



## EXAM 1\_3 (Algebraic Fractions - Equations)

1. Work out and simplify:

(2.5 points)

a)  $\left(1 + \frac{1}{x-1}\right) : \frac{x^2+x}{x^2-1} =$

b)  $\frac{x^2-3x}{x+2} : \frac{x^2-5x+6}{x^2-4} =$

2. Solve the following:

(1.5 points each equation)

a)  $\frac{x}{x-2} - \frac{2}{x-2} = \frac{6}{x}$

b)  $x + \sqrt{3x+10} = 6$

c)  $x^4 + x^3 - 4x^2 - 4x = 0$

d)  $(x^2 + 3x)^2 - 3(2x^3 + 4x^2 - 2) = 2 - 4(x-2)(x+2)$

e)  $\sqrt{2x} - \sqrt{x+1} = -1$

## SOLUTION

$$1. a) \left(1 + \frac{1}{x-1}\right) : \frac{x^2 + x}{x^2 - 1} = \left(\frac{x-1+1}{x-1}\right) : \frac{x^2 + x}{x^2 - 1} = \frac{x(x^2 - 1)}{(x-1)(x^2 + x)} =$$

$$= \frac{x(x-1)(x+1)}{(x-1) \cdot x(x+1)} = 1$$

$$b) \frac{x^2 - 3x}{x+2} : \frac{x^2 - 5x + 6}{x^2 - 4} = \frac{(x^2 - 3x)(x^2 - 4)}{(x+2)(x^2 - 5x + 6)}$$

$$= \frac{x(x-3)(x-2)(x+2)}{(x+2)(x-3)(x-2)} = x$$

$$x^2 - 3x = x(x-3) \quad x^2 - 5x + 6 = 0 \Rightarrow x = \frac{5 \pm \sqrt{25 - 24}}{2} = \begin{cases} 3 \\ 2 \end{cases}$$

$$x^2 - 4 = (x+2)(x-2) \quad x^2 - 5x + 6 = (x-3)(x-2)$$

$$2. a) \frac{x}{x-2} - \frac{2}{x-2} = \frac{6}{x} \Rightarrow \frac{x^2 - 2x}{x(x-2)} = \frac{6(x-2)}{x(x-2)} \Rightarrow x^2 - 8x + 12 = 0 \Rightarrow x = \begin{cases} 6 \\ 2 \end{cases}$$

$x = 2$  is not a solution, this value would cause division by zero in the original equation. Since the solution is  $x = 6$

$$b) x + \sqrt{3x+10} = 6 \Rightarrow \sqrt{3x+10} = 6 - x \Rightarrow 3x+10 = (6-x)^2$$

$$3x+10 = 36 + x^2 - 12x \Rightarrow 0 = x^2 - 15x + 26$$

$$x = \frac{15 \pm \sqrt{225 - 104}}{2} = \frac{15 \pm \sqrt{121}}{2} = \frac{15 \pm 11}{2} \rightarrow \begin{cases} x=13 \\ x=2 \end{cases}$$

Check:

$$x = 13 \rightarrow 13 + \sqrt{49} = 13 + 7 = 20 \neq 6 \rightarrow x = 13 \text{ no!}$$

$$x = 2 \rightarrow 2 + \sqrt{16} = 2 + 4 = 6 \rightarrow x = 2 \text{ yes!}$$

SOLUTION:  $x = 2$

c)  $x^4 + x^3 - 4x^2 - 4x = 0$ , we factorize the polynomial:

$$x^4 + x^3 - 4x^2 - 4x = x(x^3 + x^2 - 4x - 4)$$



$$\text{Div}(4) = \pm 1, \pm 2, \pm 4$$

$$P(1) = 1 + 1 - 4 - 4 \neq 0$$

$$P(-1) = -1 + 1 + 4 - 4 = 0$$

	1	+1	-4	-4	$x^2 - 4 = 0$
-1		-1	0	+4	$x = \sqrt{4} = \pm 2$
	1	0	-4	0	

$$\text{EQUATION: } x(x+1)(x-2)(x+2) = 0$$

$$\text{SOLUTION: } x = 0; x = -1; x = 2; x = -2$$

$$d) (x^2 + 3x)^2 - 3(2x^3 + 4x^2 - 2) = 2 - 4(x-2)(x+2)$$

$$x^4 + 6x^3 + 9x^2 - 6x^3 - 12x^2 + 6 = 2 - 4(x^2 - 4)$$

$$x^4 + 9x^2 - 12x^2 + 6 = 2 - 4x^2 + 16 \rightarrow x^4 + x^2 - 12 = 0 \quad \text{biquadratic}$$

$$z = x^2 \rightarrow z^2 + z - 12 = 0 \rightarrow z = \frac{-1 \pm \sqrt{1+48}}{2} = \begin{cases} 3 \\ -4 \end{cases} \rightarrow x = \begin{cases} \pm \sqrt{3} \\ \pm \sqrt{-4} \end{cases}$$

$$\text{SOLUTION: } x = \pm \sqrt{3}$$

$$e) \sqrt{2x} - \sqrt{x+1} = -1 \Rightarrow \sqrt{2x} + 1 = \sqrt{x+1} \Rightarrow (\sqrt{2x} + 1)^2 = (\sqrt{x+1})^2$$

$$(\sqrt{2x})^2 + 2\sqrt{2x} + 1 = x + 1 \Rightarrow 2\sqrt{2x} = x + 1 - 2x - 1 \Rightarrow 2\sqrt{2x} = -x$$

$$(2\sqrt{2x})^2 = (-x)^2 \Rightarrow 4 \cdot 2x = x^2 \Rightarrow 8x - x^2 = 0 \Rightarrow x(8-x) = 0 \Rightarrow \begin{cases} x = 0 \\ x = 8 \end{cases}$$

Check:

$$\sqrt{0} - \sqrt{0+1} = 0 - 1 = -1 \quad \text{YES}$$

$$\sqrt{16} - \sqrt{8+1} = 4 - 3 = 1 \neq -1 \quad \text{NO}$$

$$\text{SOLUTION } x = 0$$