

## CONTROL INTEGRALES INDEFINIDAS

$$1) \int (x^2 + 7x - 5) \cdot \cos x \, dx$$

$$2) \int x \tan^2 x \, dx$$

$$3) \int \frac{\ln^2 x}{x^2} \, dx$$

$$4) \int \frac{dx}{2\sin^2 x + 3\cos^2 x}$$

$$5) \int \frac{e^x + e^{2x} + e^{3x}}{e^{4x}} \, dx$$

$$6) \int \frac{2x^2 + x - 1}{3x(x+2)(x-2)} \, dx$$

7) Sea  $f : (-1,1) \rightarrow \mathbb{R}$  la función definida por  $f(x) = \ln(1-x^2)$ . Calcula la primitiva de  $f$  cuya gráfica pasa por el punto  $(0,1)$

## SOLUCIONES

$$\begin{aligned}
 1) \int (x^2 + 7x - 5) \cos x \, dx &= (\text{llamamos} \quad \left. \begin{array}{l} u = x^2 + 7x - 5 \\ dv = \cos x \, dx \\ v = \sin x \end{array} \right\} \quad ) = \\
 &= (x^2 + 7x - 5) \sin x - \int (2x + 7) \sin x \, dx = (\text{haciendo} \quad \left. \begin{array}{l} u = 2x + 7 \\ dv = \sin x \, dx \\ v = -\cos x \end{array} \right\} ) \\
 &= (x^2 + 7x - 5) \sin x + (2x + 7) \cos x - 2 \int \cos x \, dx = \\
 &= (x^2 + 7x - 7) \sin x + (2x + 7) \cos x + C
 \end{aligned}$$

$$\begin{aligned}
 2) \int x \tan^2 x \, dx &= (\left. \begin{array}{l} u = x \\ dv = \tan^2 x \, dx \\ v = \tan x - x \end{array} \right\} \quad ) = \\
 &(*) v = \int \frac{\sin^2 x}{\cos^2 x} \, dx = \int \frac{1 - \cos^2 x}{\cos^2 x} \, dx = \int \frac{1}{\cos^2 x} \, dx - \int 1 \, dx = \tan x - x \\
 &= x \tan x - x^2 - \int (\tan x - x) \, dx = x \tan x - \frac{x^2}{2} - \ln(\cos x) + C
 \end{aligned}$$

$$\begin{aligned}
 3) \int \frac{\ln^2 x}{x^2} \, dx &= (\left. \begin{array}{l} u = \ln^2 x \\ dv = \frac{1}{x^2} \, dx \\ v = -\frac{1}{x} \end{array} \right\} \quad ) = -\frac{1}{x} \ln^2 x + 2 \int \frac{\ln x}{x^2} \, dx = (\text{y ahora,} \\
 &\left. \begin{array}{l} u = \ln x \\ dv = \frac{1}{x^2} \, dx \end{array} \right\} \quad \left. \begin{array}{l} du = \frac{dx}{x} \\ v = -\frac{1}{x} \end{array} \right\} = -\frac{1}{x} \ln^2 x + 2 \left[ -\frac{\ln x}{x} + \int \frac{dx}{x^2} \right] = -\frac{1}{x} (\ln^2 x + 2 \ln x + 2) + C
 \end{aligned}$$

$$\begin{aligned}
 4) \int \frac{dx}{2\sin^2 x + 3\cos^2 x} &= (\text{Dividimos numerador y denominador entre } 3\cos^2 x) = \\
 &= \int \frac{\frac{1}{3\cos^2 x} \, dx}{\frac{2\sin^2 x}{3\cos^2 x} + 1} = \frac{1}{3} \int \frac{\frac{1}{\cos^2 x} \, dx}{\left( \sqrt{\frac{2}{3}} \tan x \right)^2 + 1} = (\left. \begin{array}{l} t = \sqrt{\frac{2}{3}} \tan x \\ dt = \sqrt{\frac{2}{3}} \frac{1}{\cos^2 x} \, dx \end{array} \right\} ) = \frac{1}{\sqrt{6}} \operatorname{arctg} \left( \sqrt{\frac{2}{3}} \tan x \right) + C
 \end{aligned}$$

$$\begin{aligned}
 5) \int \frac{e^x + e^{2x} + e^{3x}}{e^{4x}} \, dx &= \int e^{-3x} \, dx + \int e^{-2x} \, dx + \int e^{-x} \, dx = \\
 &= -\frac{1}{3} e^{-3x} - \frac{1}{2} e^{-2x} - e^{-x} + C
 \end{aligned}$$

$$6) \int \frac{2x^2 + x - 1}{3x(x+2)(x-2)} dx = \int \frac{1/4}{3x} dx + \int \frac{5/24}{x+2} dx + \int \frac{9/24}{x-2} dx =$$

$$\frac{1}{12} \ln|x| + \frac{5}{24} \ln|x+2| + \frac{9}{24} \ln|x-2| + C$$

7) Empezamos calculando  $\int \ln(1-x^2)dx$  que se hace por partes:

$$\begin{aligned} u &= \ln(1-x^2) \\ dv &= dx \\ \left. \begin{array}{l} du = \frac{-2x}{1-x^2} dx \\ v = x \end{array} \right\} \Rightarrow \int \ln(1-x^2)dx = x \ln(1-x^2) - \int \frac{-2x^2}{1-x^2} dx = \\ &= x \ln(1-x^2) + 2 \int \frac{x^2}{1-x^2} dx \text{ para hacer esta integral, dividimos:} \\ &\frac{x^2}{-x^2+1} \quad \left| \begin{array}{c} -x^2+1 \\ -1 \end{array} \right. \quad \Rightarrow \frac{x^2}{1-x^2} = \frac{-1(1-x^2)+1}{1-x^2} = -1 + \frac{1}{1-x^2} \end{aligned}$$

$$\text{Con lo que tenemos } \int \frac{x^2}{1-x^2} dx = \int -1 dx + \int \frac{1}{1-x^2} dx = -x + \int \frac{A}{1-x} dx + \int \frac{B}{1+x} dx$$

$$\frac{1}{1-x^2} = \frac{A}{1-x} + \frac{B}{1+x} \Rightarrow 1 = A(1+x) + B(1-x), \text{ sustituimos para } x=1 \text{ y } x=-1 \text{ y}$$

tenemos que  $A = \frac{1}{2}$  y  $B = \frac{1}{2}$ , de donde, nuestra integral, nos quedará:

$$\int \ln(1-x^2)dx = x \ln(1-x^2) + 2 \left[ -x + \frac{1}{2} \ln|1-x| + \frac{1}{2} \ln|1+x| \right] + C, \text{ es decir:}$$

$\int \ln(1-x^2)dx = x \ln(1-x^2) - 2x + \ln|1-x^2| + C$  y, como la gráfica de la primitiva pedida pasa por  $(0,1)$ , tendremos que:  $0 - 0 + \ln 1 + C = 1 \Rightarrow C = 1$  por lo que la primitiva pedida es  $x \ln(1-x^2) - 2x + \ln|1-x^2| + 1$